Readigraft® BLX DBM & CC Putty



ReadiGRAFT® BLX DBM & CC Putty is a new addition to the ReadiGRAFT BLX DBM family. It is an optimally demineralized, sterile, DBM Putty combined with cortical cancellous and uses a natrual human tissue carrier.

Brand

ReadiGRAFT® BLX DBM & CC Putty

Clinical

Any surgical application which requires bone void filler

Applications

- Optimal Osteoinductive Potential: All cortical particulate used in ReadiGraft BLX DBM & CC Putty are demineralized to a residual calcium level, proven to provide the highest osteoinductive potential.1,2,3,4,5
- Natural Osteoinductive Scaffold: The natural porous architecture of cancellous provides an optimal scaffold for cellular attachment and vascular growth.
- **Features** And Benefits
- **Excellent Handling Properties:** It can be manipulated into any deisred shape and will not migrate under irrigation.
- **Moldable:** Allows surgeon to shape product to meet intra-operative needs.
- Sterile: Sterilized using proprietary and patented Allowash XG® technology. This technology provides a sterility assurance level of 10⁻⁶, without compromising the grafts inherent osteoconductivity or osteoinductive potential.
- **Ready to Use:** No need to waste valuable OR time preparing the product.
- Natural Human Tissue Carrier: Biocompatible, resorbs naturally during bone healing.
- Innovative Syringe Design: Controlled extrusion of all available product.


Alternative **Products**

- ReadiGraft BLX Sponge
- ReadiGraft® BLX Putty
- ReadiGraft® BLX Fibers
- FlexiGraft® DBM Fibers



Readigraft® BLX DBM & CC Putty





1864 Concert Drive Virginia Beach, VA 23453 1-757-464-4761 x2000

www.AccessLifeNetHealth.org

The LifeNet Health, ReadiGraft and FlexiGraft logos are registered trademarks of LifeNet Health. ©2012 LifeNet Health, Virginia Beach, VA. All rights reserved.

68-40-102-02. 00



- 1. Effect(s) of the demineralization process on the osteoinductivity of demineralized bone matrix Zhang M, Powers R M, and Wolfinberger L. Periodontol 1997: 68:1085-1092
- 2. The affects of residual calcium in decalcified freeze-dried bone allograft in a critical-sized defect in the Rattus norvegicus calvarium Turonis JW, McPherson JC 3rd, Cuening MF: J Oral Implantol. 2006; 32(2):55-62
- 3. Effects of Varying degrees of Allograft Decalcification on the Cultured Porcine Osteoclast cells Herold RW, Pashley DH, Cuening MF: J Periodontal. 2002 Feb; 73(2): 213-9
- 4. Enhancement of osteoblast proliferation in vitro by selective enrichment of demineralized freeze-dried bone allograft with specific growth factors Mott DA., Maihot J, Cuenin MF, Sharaway M, Borke J; J Oral Implantol 2002; 28(2): 57-66
- 5. BMP depletion occurs during prolonged acid demineralization of bone: characterization and implications for graft preparation Pietrzak WS, Ali SN, Chitturi D, Jacob M, Woodell-May JE. Cell Tiss. Bank 2007 (Published on line)

International Use Only